

RASPBERRY

Take first fully expanded leaf out from the crown of the plant. The trifoliate leaf is formed of three smaller leaflets - **BE SURE TO TAKE WHOLE LEAF AND STEM (petiole), not just the leaflets.**

Take at least **140 trifoliate leaves**, randomly across the sampling area. The part to be tested is the petiole, or leaf stalk.

If you require a Sap 4 comprehensive analysis please collect **twice** as many leaves.

Monitoring program

Begin sampling when first new leaves reach mature stage. Sample fortnightly until the end of the main harvest period.



SEND PETIOLE

RASPBERRY

DIFFERENTIAL SAP ANALYSIS



Take first fully expanded leaf out from the crown of the plant. The trifoliate leaf is formed of three smaller leaflets - BE SURE TO TAKE WHOLE LEAF AND STEM (petiole), not just the leaflets. This will become your NEW LEAF SAMPLE.

Take fully expanded leaf from old part of the plant. The trifoliate leaf is formed of three smaller leaflets - BE SURE TO TAKE WHOLE LEAF AND STEM (petiole), not just the leaflets. This will become your OLD LEAF SAMPLE

Monitoring program

Begin sampling when first new leaves reach mature stage. Sample fortnightly until the end of the main harvest period.



NEW LEAF

(1ST FULLY EXPANDED LEAF)

OLD LEAF